

Purina® Goat Nutrition Program

Purina[®] Dairy Parlor 16

Supplement for lactating dairy goats

Dairy Parlor 16 feed is a pelleted feed formulated for optimum milk production in lactating dairy goats.

Purina® products deliver the nutrition and performance you expect. Try Purina® Dairy Parlor 16 feedand see the difference it can make in your goats' overall performance and health.



FEATURE	BENEFIT
Highly Palatable	Formulated specifically for dairy goats
Nutritionally Balanced	Formulated for optimal milk production and to support fertility, healthy skin and coat and overall health
Palatable, Premium Ingredients	Consistent quality helps assure high palatability and supports performance
Diamond V® Yeast Culture	Helps support feed digestion and helps to stabilize rumen fermentation during stress
Availa-4® Minerals	Balanced combination of organic zinc, manganese, copper and cobalt to optimize growth and help support immune response
Pelleted	No separation of ingredients, easy to handle





Goat

Purina® Goat Nutrition Program

Purina[®] Dairy Parlor 16



Guaranteed Analysis:

NUTRIENT	LEVEL
Crude Protein, (min)	16.00%
Crude Fat, (min)	3.00%
Crude Fiber, (max)	8.00%
Acid Detergent Fiber (ADF)	11.00%
Calcium (Ca), (min)	0.70%
Calcium (Ca), (max)	1.20%
Phosphorus (P), (min)	0.60%
Salt (NaCl), (min)	0.25%
Salt (NaCl), (max)	0.75%
Copper (Cu), (min)	20.00 ppm
Copper (Cu), (max)	25.00 ppm
Selenium (Se), (min)	0.60 ppm
Vitamin A, (min)	7,500 IU/lb
Vitamin D ₃ , (min)	750 IU/lb
Vitamin E, (min)	30 IU/lb

Ingredients:

Processed Grain By-Products, Grain Products, Plant Protein Products, Roughage Products, Molasses Products, Calcium Carbonate, Salt, Sodium Sesquicarbonate, Soybean Oil, Niacin Supplement, Zinc Sulfate, Vitamin E Supplement, Vitamin A Supplement, Manganese Sulfate, Zinc Oxide, Manganous Oxide, Zinc Amino Acid Complex, Sodium Selenite, Manganese Amino Acid Complex, Cobalt Carbonate, Copper Amino Acid Complex, Vitamin D₃ Supplement, Ferrous Sulfate, Ethylenediamine Dihydroiodide, Cobalt Glucoheptonate, Potassium Iodide, Basic Copper Chloride, Sodium Molybdate.

This product is available regionally, so please check with your local Purina® dealer for ordering details.

Directions:

Feed this product according to production requirements (1 pound of lactation ration per every 2 quarts of milk produced) along with good quality forage free-choice. Do not dilute with grain. Feed intake will vary with forage quality, life stage, environment and activity level.

Important:

- 1. Start kids on colostrum milk from birth to 3 days of age; feed Purina® Kid Milk Replacer from 2 days until 4-8 weeks of age.
- If kids have not been creep fed or they were purchased, then they should be placed on grass hay on a free-choice basis after arrival.
- 3. When making a ration change, allow 3 to 5 days for animals to adjust to the new ration.
- Keep a constant supply of complete feed available. Do not let fine material accumulate in feeders
- Provide adequate bunk space for each animal. Bunks should be well-protected and well-managed to prevent feed from becoming wet or moldy.
- 6. Provide a source of fresh, clean water at all times.
- 7. Feed Purina Goat Mineral free-choice.
- 8. Consult your veterinarian for the recommended health program for your local area. This includes internal and external parasite control. In groups of goats there are certain animals that experience chronic bloat, or other digestive disturbances and consequently, are poor performers. In addition, excess feed consumption, severe weather changes resulting in erratic feed consumption and poorly managed feeding practices, can increase the incidence of bloat in all goats. If bloating occurs, the above management practices should be reviewed and your veterinarian consulted.

Caution:

Store in a dry, well-ventilated area protected from rodents and insects. Do not feed moldy or insect-infested feed to animals as it may cause illness, performance loss or death.

This product contains copper and should not be fed to sheep.

USE ONLY AS DIRECTED.

